

Published in Radiation Res. 43:152-160, 1970

AFRRI SR69-24

DECEMBER 1969

AFRRI
SCIENTIFIC
REPORT

FOR REFERENCE

Do Not Take From This Room

**EFFECT OF IONIZING RADIATIONS
ON DISTRIBUTION OF PLASMA
PROTEIN-BOUND NEUTRAL HEXOSES
IN MICE AND DOGS**

AFRRI SR69-24

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
Defense Atomic Support Agency
Bethesda, Maryland

Distribution of this document is unlimited.

This report has been approved for open publication by the Department of Defense

All aspects of investigative programs involving the use of laboratory animals sponsored by DOD components are conducted according to the principles enunciated in the "Guide for Laboratory Animal Facilities and Care", prepared by the National Academy of Sciences - National Research Council.


December 1969

EFFECT OF IONIZING RADIATIONS ON DISTRIBUTION OF PLASMA
PROTEIN-BOUND NEUTRAL HEXOSES IN MICE AND DOGS

A. S. EVANS



R. E. GEORGE
Commander, MSC, USN
Chairman
Radiation Biology Department



HUGH B. MITCHELL
Colonel, USAF, MC
Director

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE
Defense Atomic Support Agency
Bethesda, Maryland

Distribution of this document is unlimited

TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary)	iii
Abstract	vii
I. Introduction	1
II. Materials and Methods	1
III. Results	3
IV. Discussion	8
References	11

LIST OF FIGURES

	Page
Figure 1. Percent increase in concentration of total neutral hexoses in the plasma of dogs at indicated times after exposure to mixed gamma-neutron radiations	3
Figure 2. Milligrams protein-bound neutral hexoses per 100 mg protein (biuret) in the plasma of dogs at indicated times relative to exposure to mixed gamma-neutron radiations	4
Figure 3. Typical acrylamide gel disc electrophoretic patterns of plasma of a dog and of mice at indicated intervals relative to lethal irradiation with mixed gamma-neutron radiations	5
Figure 4. Characteristic protein absorption curve (280 nm) of plasma from mice and dogs illustrating the four fractions regularly obtained by combined molecular sieving and ion exchange chromatography	6
Figure 5. Protein-bound neutral hexoses in fractions from combined molecular sieving and ion exchange chromatography columns	7
Figure 6. Acrylamide gel disc electrophoretic patterns of material from fractions I and III of Figure 5 (dog)	8

FOREWORD

(Nontechnical summary)

Practically all of the proteins in the blood plasma are complex molecules containing carbohydrates (sugars) in addition to amino acids as a part of their primary structure.

In the first report of this series (AFRRI SR 68-4), it was suggested that blood plasma concentration levels of protein-bound carbohydrates (PBC) as neutral hexoses (uncharged 6-carbon sugars) may constitute a crude index of radiosensitivity prior to and prognosis after irradiation in otherwise healthy C3H mice.

The presently reported experiments were designed to test whether dogs would exhibit radiation-induced PBC response in plasma similar to that of mice and to identify operationally the specifically affected plasma protein components in both species.

C3H mice and beagles were subjected to single, whole-body exposures of mixed gamma-neutron radiations delivered as a pulse or at a steady-state rate of approximately 20 rads/min. Blood specimens were taken before irradiation and at daily intervals for varying periods, up to 20 days. PBC concentration was determined by chemical analysis.

The affected protein components were identified operationally by sieving through a gel which had a pore size small enough to separate the various proteins roughly by molecular size and at the same time contained groups which accelerated or retarded the molecules according to their electrical charge. The fractions from these columns were then examined by separating the proteins in each by their relative movement in a molecular sieving polyacrylamide gel under the influence of an electrical field.

Further investigations to test the hypothesis that the preirradiation PBC concentration may constitute a crude index to radiosensitivity of the individual were made under specifically designed conditions. The mice and dogs were bled prior to irradiation only. The plasma PBC values of each group were arranged in three sets by a standard statistical procedure: those which were in a zone immediately about the group mean, those below the lower limit of that zone, and those above the upper limit. The hypothesis being tested postulated that the animals with plasma PBC concentrations in the low set would be relatively radioresistant, those in the high set would be relatively radiosensitive, and the set within the limits of the mean value would be undistributed with regard to mortality.

Alterations in plasma concentration of protein-bound carbohydrates followed the same general course in dogs as previously reported for mice. Thus, in the animals which died, the PBC concentration showed a marked increase, while the survivors deviated only slightly from their preirradiation base-line values. The similarity of the radiation-induced plasma PBC response of mice and dogs suggested that this phenomenon may be generally applicable to mammals and not species specific.

A large portion of the increase in PBC in animals near death was accounted for operationally in carbohydrate containing proteins which normally function to help conserve the body's supply of iron.

The findings of these experiments were interpreted to reflect radiation-induced damage to bodily control mechanisms which resulted in excessive circulating concentrations of these proteins. The stability of the plasma PBC concentration maintained

by the survivors suggested that their bodily control mechanisms had sufficient reserve to enable them to maintain balance despite the radiation insult.

The usefulness of the preirradiation PBC concentration as an indicator of radiosusceptibility was confirmed.

The results reported here indicate that changes in plasma concentration of PBC may be developed into a relatively simple, objective prognostic test to supplement clinical observation in cases of radiation injury.

ABSTRACT

Comparison of changes in concentration of protein-bound carbohydrates (PBC) as neutral hexoses in the plasma of C3H mice and beagles as a function of time after exposure to mixed gamma-neutron radiations revealed that the two species react similarly. The animals which died exhibited a marked increase in plasma PBC concentration while the survivors deviated only slightly from their preirradiation values. Analytical acrylamide gel electrophoresis demonstrated that distinctive, progressive postirradiation changes occurred in the plasma of the animals which succumbed. The most striking changes were in the components tentatively identified as transferrin, haptoglobins, and α_2 -macroglobulin. Preparative fractionation of plasma of both mice and dogs by combined molecular sieving and ion exchange chromatography revealed that, in the four fractions uniformly obtained, absolute changes in PBC regularly occurred only in the first and third. The protein composition of these fractions and their significance as a prognostic tool in radiation injury are discussed. The suggestion that the preirradiation PBC concentration may constitute a crude index to radiosensitivity was confirmed under conditions specifically designed to test the hypothesis.

I. INTRODUCTION

Evans et al.⁴ suggested that plasma concentration levels of protein-bound carbohydrates (PBC) as neutral hexoses may constitute a crude index of radiosensitivity prior to and prognosis after irradiation in otherwise healthy C3H mice. They pointed out, however, that the glycoproteins constitute such a structurally and physiologically diverse agglomeration that analyses which treat them as a single entity cannot be expected to yield clinically applicable criteria.

The presently reported experiments were designed to test whether dogs would exhibit radiation-induced PBC response in plasma similar to that of mice, to identify operationally the specifically affected plasma protein components, and to test the hypothesis that the preirradiation PBC concentration may constitute a crude index to radiosensitivity of the individual.

II. MATERIALS AND METHODS

Two species of animals were utilized in these experiments: young adult male C3H mice, 6-8 weeks of age, weighing between 20 and 24 g, and healthy AKC registrable beagles of both sexes, 2-3 years of age.

Blood specimens were taken before irradiation and at daily intervals for varying periods, up to 20 days. Some of the dogs were utilized in cooperation with other investigators so that doses, dose rates, and intervals of blood sampling were those which would not interfere with the primary experimental design.

All of the animals were exposed to mixed gamma-neutron radiations from the AFRRI-TRIGA reactor. The characteristics of the exposure field and the array for exposure of the mice have been described previously.⁴ The dogs were exposed

unilaterally to either pulsed or steady-state doses, the details of the methodology of which have been described.⁷ The midline tissue doses (rads) and dose rates (rads/min) received by the animals reported on here are given with the analytical data.

Protein-bound neutral hexoses (as galactose and mannose) were estimated by a modification⁴ of the sulfuric acid-orcinol method of Weimer and Moshin.⁹

Combined molecular sieving and ion exchange chromatography were accomplished utilizing a porous polyacrylamide gel with a functional cation exchange group ($\dots\text{COO}^- \text{Na}^+$) attached to the gel matrix.* The fractions were eluted from the 0.9 x 12 cm columns by stepwise increases in ionic strength and pH of 0.05 M phosphate (Na^+) buffers as indicated with the experimental data.

It was necessary to pool plasma from similarly reacting mice to obtain sufficient amounts for the columns (0.5 ml). The size of the dogs permitted withdrawal of sufficient blood to allow analyses on individual animals.

The distribution of protein-bound neutral hexoses was determined by precipitation of each of the 0.5 ml collections from the chromatography column with an equal volume of 10 percent aqueous trichloroacetic acid. The precipitate was washed with 95 percent ethanol, redissolved in 0.1 N NaOH, and the sulfuric acid-orcinol procedure applied as for whole plasma.⁹

The protein composition of each of the fractions from the chromatography columns was examined by analytical polyacrylamide gel electrophoresis carried out essentially as described by Davis³ and stained with either Buffalo black or Coomassie blue.²

* BioGel, CM-100, Calbiochem, Los Angeles, California

To test the hypothesis that the preirradiation PBC concentration may constitute a crude index to radiosensitivity of the individual, two groups of animals, 18 dogs and 24 mice, were bled prior to irradiation only. The dogs were exposed to a 205-rad pulse ($LD_{44/30}$); the mice to 400 rads at 20 rads/min ($LD_{46/30}$).

The plasma PBC values of each group were arranged in three sets: those which were within the 95 percent confidence interval for the group mean, those below the lower limit of that interval, and those above the upper limit.

III. RESULTS

Alterations in the plasma concentration of protein-bound carbohydrates (PBC) as neutral hexoses followed the same general course in dogs (Figure 1) as previously reported for mice.⁴ Thus, in the animals which died, the PBC concentration showed a marked increase while the survivors deviated only slightly from their preirradiation base-line values.




Figure 1. Percent increase in concentration of total neutral hexoses in the plasma of dogs at indicated times after exposure to mixed gamma-neutron radiations. Numbers in parentheses are number of animals represented by each curve.

Hemodilution and hemoconcentration were eliminated as controlling factors in the changes demonstrated by expressing the neutral hexose concentration as milligrams of carbohydrate per 100 mg biuret protein (Figure 2).

Analytical acrylamide gel electrophoresis of plasma of both mice and dogs (Figure 3) revealed that distinctive, progressive postirradiation changes occurred




Figure 2. Milligrams protein-bound neutral hexoses per 100 mg protein (biuret) in the plasma of dogs at indicated times relative to exposure to mixed gamma-neutron radiations. The doses were 225 rads: ○ = 30-day survivors; ● = died on 16th day postirradiation; 230 rads: ▼ = died on 12th day; ▲ = died on 15th day; 400 rads: ◆ = died on 4th day; □ = died on 10th day; ■ = died on 11th day. Numbers in parentheses are number of animals represented by each curve.

in the animals which succumbed. The most striking changes appeared in components identified as transferrin, haptoglobin and α_2 -macroglobulin.




Figure 3. Typical acrylamide gel disc electrophoretic patterns of plasma of a dog (A), and of mice (B), at indicated intervals relative to lethal irradiation with mixed gamma-neutron radiations.

Preparative fractionation of plasma of both dogs and mice by combined molecular sieving and ion exchange chromatography regularly gave four protein fractions as measured by monitoring the column at 280 nm wavelength (Figure 4). A marked




Figure 4. Characteristic protein absorption curve (280 nm) of plasma from mice (upper) and dogs (lower) illustrating the four fractions regularly obtained by combined molecular sieving and ion exchange chromatography. Note marked postirradiation increase in fraction III. Analyses of the four fractions are given in Figure 5.

increase in absorbance at this wavelength was seen in fraction III in all the patterns of moribund animals.

Chemical analyses revealed that all the radiation-induced increases in plasma PBC content were in the first and third fractions (Figure 5).




Figure 5. Protein-bound neutral hexoses in fractions from combined molecular sieving and ion exchange chromatography columns. For designation of fraction numbers see Figure 4.

Analytical acrylamide gel disc electrophoresis demonstrated that the increase in PBC regularly found in the first fraction was attributable to the increase in transferrin (compare Figure 6a and 6b). The third fraction in which the absolute PBC content increased up to more than eight times the preirradiation value, contained the haptoglobin complex, β_2 -glycoproteins, and the α_2 -macroglobulin (Figure 6c and 6d).

Of the 18 dogs on which only preirradiation blood samples were taken, 8 died during the 30-day observation period and 10 survived. The 95 percent confidence interval of the mean of 121.2 mg PBC/100 ml plasma was 112.4 to 130.1. All of the

five animals which had PBC values below the lower limit survived; four of the six which were above the upper limit on the mean died.

In the 24 mice, 11 of which died, the confidence interval of the mean of 159.1 mg PBC/100 ml plasma was 154.3 to 163.8. Eight of the nine mice which had PBC values below the lower limit survived, while eight of the ten above the upper limit died.




Figure 6. Acrylamide gel disc electrophoretic patterns of material from fractions I and III of Figure 5 (dog): (a) fraction I, preirradiation; (b) fraction I, 7 days postirradiation; (c) fraction III, preirradiation; (d) fraction III, 7 days postirradiation. Note the marked increase in transferrin (\Rightarrow).

IV. DISCUSSION

The similarity of the radiation-induced plasma PBC response of mice and dogs as a function of time after exposure suggested that this phenomenon may be generally applicable to mammals and not species specific.

It was apparent that a large portion of the increase in protein-bound neutral hexoses in moribund animals could be accounted for operationally (Figures 3 and 6) in glycoproteins concerned with iron conservation, transferrin and haptoglobin. It is doubtful that the marked increase in these plasma components is totally a response to a biological need for their primary functions, although compensatory attempts to bolster declining erythropoiesis by conserving iron and hemoglobin may have been a factor.

Considering the rapidity of the rise, especially in the animals which died in the first week, it did not appear likely that the elevation in transferrin was as a compensatory defense mechanism.⁶ The consistency of this increase compared with the variability in loss of γ -globulins⁵ did not support such a hypothesis.

More likely, the findings reported here reflect radiation-induced damage to control mechanisms of biosynthesis and catabolism resulting in excessive circulating concentrations of these proteins. That the surviving animals did not show significant changes lends credence to this hypothesis. Thus, the stability of plasma concentration of protein-bound neutral hexoses maintained by the survivors indicated that their metabolic control mechanisms had sufficient reserve to enable them to maintain physiological balance despite the radiation insult.

The marked rise in the α_2 -macroglobulin component in the moribund animals explains the great increase over preirradiation levels in absorbance of the plasma at 280 nm wavelength in fraction III of Figure 4. This protein has been shown⁸ to contain over 500 residues per molecule of aromatic amino acids which absorb at or near 280 nm.

The carbohydrate composition of the proteins which have been shown here to account for most if not all of the rise in plasma PBC concentration in moribund animals is similar in the relative amounts of the various classes of sugars they contain. It is expected, therefore, that further analyses should show the neutral hexoses, hexosamines, and sialic acid will remain in direct proportion to one another during the course of radiation sickness. Thus, differentiation of radiation injury from hypertension, myocardial infarction, and liver disorders¹ will be permitted. Experiments are in progress to test this hypothesis.

The results reported here indicate that changes in concentration of circulating protein-bound neutral hexoses may be developed into a relatively simple, objective prognostic test to supplement clinical observation in cases of radiation injury.

The confirmation, under conditions specifically designed to test the thesis that preirradiation levels of protein-bound neutral hexoses permit screening of small groups for extremes in radiosusceptibility offers promise. The general concept has been consistent in this⁴ and other¹⁰ laboratories but has been restricted to studies using inbred strains of mice, rats, and dogs. Tests of its applicability to the mongrel human species present obvious difficulties in experimental design.

REFERENCES

1. Anantha Samy, T. S., Patel, S. J. and Cama, H. R. Significance of serum protein-bound carbohydrates in diagnosis of hypertension, myocardial infarction and liver disorders. *Indian J. Med. Res.* 55:169-173, 1967.
2. Chrambach, A., Reisfeld, R. A., Wyckoff, M. and Zaccari, J. A procedure for rapid and sensitive staining of protein fractionated by polyacrylamide gel electrophoresis. *Anal. Biochem.* 20:150-154, 1967.
3. Davis, B. J. Disc electrophoresis -- II. Method and application to human serum proteins. *Ann. N. Y. Acad. Sci.* 121:404-427, 1964.
4. Evans, A. S., Quinn, F. A., Brown, J. A. and Strike, T. A. Effect of ionizing radiation on total protein-bound neutral hexoses in the plasma of mice. *Radiation Res.* 36:128-137, 1968.
5. Evans, A. S., Quinn, F. A. and Strike, T. A. Effect of monoenergetic 14-MeV neutron and mixed gamma-neutron radiations on immunoelectrophoretic patterns of mice. *Radiation Res.* 34:49-55, 1968.
6. Martin, C. M. Relation of transferrin to serum bacteriostatic activity in agammaglobulinemic and other patients. *Am. J. Med. Sci.* 244:334-336, 1962.
7. Pitchford, T. L. and Thorp, J. W. The acute mortality response of beagles to pulsed, mixed gamma-neutron radiations. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-15, 1968.
8. Schultze, H. E. and Heremans, J. F. *Molecular Biology of Human Proteins with Special Reference to Plasma Proteins*, Vol. 1. Amsterdam, London, New York, Elsevier Publishing Company, 1966.
9. Weimer, H. E. and Moshin, J. R. Serum glycoprotein concentrations in experimental tuberculosis of guinea pigs. *Am. Rev. Tuberculosis* 68:594-602, 1953.
10. Zicha, B. O povaze změn sérových bílkovin v periferní krvi po celotělovém ozaření, pp. 75-84. In: *Experimentální postiradiační syndrom*, Praha, Czechoslovakia, Státní zdravotnické nakladatelství, 1966. *Nucl. Sci. Abstr.* 22:5297 (Abstract 52004), 1968.

DISTRIBUTION LIST

AIR FORCE

The Surgeon General, U. S. Department of the Air Force, Washington, D. C. 20333 (1)
Executive Officer, Director of Professional Services, Office of the Surgeon General, Hq. USAF (AFMSPA) T-8,
Washington, D. C. 20333 (1)
Headquarters, U. S. Air Force (AFMSPAB), Washington, D. C. 20333 (1)
Chief, Weapons and Weapons Effects Division, Hq. RTD (RTTW), Bolling AFB, Washington, D. C. 20332 (1)
Office of the Command Surgeon (ADCSC), Hq. ADC, USAF, Ent AFB, Colorado 80912 (1)
Commander, 6571st Aeromedical Research Laboratory, Holloman AFB, New Mexico 88330 (2)
Air Force Weapons Laboratory, ATTN: WLIL (1), ATTN: WLRB-2 (1), Kirtland AFB, New Mexico 87117 (2)
Chief, Nuclear Medicine Department, P. O. Box 5088, USAF Hospital Wright-Patterson, Wright-Patterson AFB,
Ohio 45433 (1)
USAFSAM (SMBR), ATTN: Chief, Radiobiology Branch, Brooks AFB, Texas 78235 (1)

ARMY

The Surgeon General, U. S. Department of the Army, Washington, D. C. 20315 (1)
Surgeon General, ATTN: MEDDH-N, U. S. Department of the Army, Washington, D. C. 20315 (1)
USACDC CSSG, Doctrine Division, Fort Lee, Virginia 23801 (1)
Commanding Officer, USACDC CBR Agency, Fort McClellan, Alabama 36201 (1)
Commanding Officer, U. S. Army Combat Developments Command, Institute of Nuclear Studies, Fort Bliss, Texas
79916 (1)
CG, USCONARC, ATTN: ATUTR-TNG (NBC), Fort Monroe, Virginia 23351 (1)
Commanding Officer, Harry Diamond Laboratories, ATTN: Nuclear Vulnerability Branch, Washington, D. C.
20438 (1)
Nuclear Branch AMCRD-DN-RE, U. S. Army Materiel Command, Washington, D. C. 20315 (1)
Commanding Officer, U. S. Army Medical Research Laboratory, Fort Knox, Kentucky 40121 (1)
Commanding Officer, USA Nuclear Medical Research Detachment, Europe, APO New York, New York 09180 (2)
Chief of Research and Development, ATTN: Nuclear, Chemical and Biological Division, U. S. Department of the
Army, Washington, D. C. 20310 (1)
Army Research Office, ATTN: Chief, Scientific Analysis Branch, Life Sciences Division, 3045 Columbia Pike,
Arlington, Virginia 22204 (1)
Division of Nuclear Medicine, Walter Reed Army Institute of Research, Walter Reed Army Medical Center,
Washington, D. C. 20012 (5)
Commanding Officer, U. S. Army Environmental Hygiene Agency, ATTN: USAEHA-RP, Edgewood Arsenal,
Maryland 21010 (1)
Commandant, U. S. Army Medical Field Service School, ATTN: MEDEW - ZNW, Fort Sam Houston, Texas
78234 (1)

NAVY

Chief, Bureau of Medicine and Surgery, U. S. Navy Department, Washington, D. C. 20390 (1)
Chief, Bureau of Medicine and Surgery, ATTN: Code 71, U. S. Navy Department, Washington, D. C. 20390 (1)
Commanding Officer, Naval Aerospace Medical Institute, Naval Aviation Medical Center, ATTN: Director of
Research, Pensacola, Florida 32512 (3)
Commanding Officer, Nuclear Weapons Training Center, Atlantic, Nuclear Warfare Department, Norfolk, Virginia
23511 (1)
Commanding Officer, Nuclear Weapons Training Center, Pacific, U. S. Naval Air Station, North Island, San Diego,
California 92135 (1)
Director, Biological Sciences Division, Office of Naval Research, Washington, D. C. 20360 (1)
Commanding Officer, U. S. Naval Hospital, ATTN: Director, REEL, National Naval Medical Center, Bethesda,
Maryland 20014 (1)
Head, Animal Behavioral Sciences Branch, Naval Aerospace Medical Institute, Naval Aerospace Medical Center,
Pensacola, Florida 32512, ATTN: Dr. John S. Thach, Jr. (1)

D.O.D.

Director, Defense Atomic Support Agency, Washington, D. C. 20305 (1)
Director, Defense Atomic Support Agency, ATTN: DDST, Washington, D. C. 20305 (1)
Director, Defense Atomic Support Agency, ATTN: Chief, Medical Directorate, Washington, D. C. 20305 (4)
Director, Defense Atomic Support Agency, ATTN: Chief, Radiation Directorate, Washington, D. C. 20305 (1)
Director, Defense Atomic Support Agency, ATTN: Technical Library, Washington, D. C. 20305 (2)
Commander, Field Command, Defense Atomic Support Agency, ATTN: FC Technical Library, Sandia Base,
Albuquerque, New Mexico 87115 (1)
Commander, Headquarters Field Command, Defense Atomic Support Agency, ATTN: FCTG8, Sandia Base,
Albuquerque, New Mexico 87115 (2)
Director, Armed Forces Institute of Pathology, Washington, D. C. 20305 (1)
Administrator, Defense Documentation Center, Cameron Station, Bldg. 5, Alexandria, Virginia 22314 (20)

OTHER GOVERNMENT

U. S. Atomic Energy Commission, Headquarters Library, Reports Section, Mail Station G-17, Washington, D. C.
20545 (1)
U. S. Atomic Energy Commission, Division of Biology and Medicine, Washington, D. C. 20545 (1)
U. S. Atomic Energy Commission, Bethesda Technical Library, 7920 Norfolk Avenue, Bethesda, Maryland 20014 (1)
National Aeronautics and Space Administration, ATTN: Lt. Col. Charles M. Barnes, USAF, DB-3, MSC, Houston,
Texas 77058 (1)
National Bureau of Standards, ATTN: Chief, Radiation Physics Division, Washington, D. C. 20234 (1)
U. S. Public Health Service, Deputy Chief, Division of Radiological Health, Washington, D. C. 20201 (1)
U. S. Public Health Service, Radiological Health Laboratory, ATTN: Library, 1901 Chapman Avenue, Rockville,
Maryland 20852 (1)
U. S. Public Health Service, Northeastern Radiological Health Laboratory, 109 Holton Street, Winchester,
Massachusetts 01890 (1)
U. S. Public Health Service, Southwestern Radiological Health Laboratory, P. O. Box 684, Las Vegas, Nevada
89101 (1)
U. S. Public Health Service, National Center for Radiological Health, Information Office, Room 3, Twinbrook
Laboratory, RBE Program, 1901 Chapman Avenue, Rockville, Maryland 20852 (1)

OTHER

Argonne National Laboratory, Library Services Department, Report Section Bldg. 203, RM-CE-125, 9700 South
Cass Avenue, Argonne, Illinois 60440 (1)
Dr. Donald G. Baker, Radiobiology Department, Zellerbach Saroni Tumor Institute, 1600 Divisadero Street,
San Francisco, California 94115 (1)
Brookhaven National Laboratory, Information Division, ATTN: Research Library, Upton, Long Island, New York
11973 (2)
Dr. J. S. Burkle, Director of Nuclear Medicine, York Hospital, York, Pennsylvania 17403 (1)
S. C. Bushong, Department of Radiology, Baylor University College of Medicine, Houston, Texas 77024 (1)
University of California, Lawrence Radiation Laboratory, Library, Bldg. 50, Room 134, Berkeley, Calif. 94720 (1)
Director, Radiobiology Laboratory, University of California, Davis, California 95616 (1)
University of California, Lawrence Radiation Laboratory, Technical Information Division Library L-3, P. O.
Box 808, Livermore, California 94551 (2)
University of California, Laboratory of Nuclear Medicine and Radiation Biology, Library, 900 Veteran Avenue,
Los Angeles, California 90024 (1)
Dr. C. Jelleff Carr, Director, Life Sciences Research Office, Federation of American Societies for Experimental
Biology, 9650 Rockville Pike, Bethesda, Maryland 20014 (1)
Director, Collaborative Radiological Health Laboratory, Colorado State University, Fort Collins, Colorado 80521 (1)
Dr. L. W. Davis, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pa. 19104 (1)
Professor Merrill Eisenbud, New York University, Tuxedo, New York 10987 (1)
Dr. T. C. Evans, Radiation Research Laboratory, College of Medicine, University of Iowa, Iowa City, Iowa 52240 (1)
Dr. Arnold Feldman, Institute of Radiology, School of Medicine, Washington University, 510 South Kingshighway,
St. Louis, Missouri 63110 (1)
Mr. Orin Gelderloos, Department of Biological Sciences, Northwestern University, Evanston, Illinois 60201 (1)
General Dynamics/Fort Worth, ATTN: Librarian, P. O. Box 748, Fort Worth, Texas 76101 (1)
Gulf General Atomic Incorporated, ATTN: Library, P. O. Box 608, San Diego, California 92112 (1)
Hazleton Nuclear Science Corporation, ATTN: Library, 4062 Fabian Way, Palo Alto, California 94303 (1)
IIT Research Institute, ATTN: Document Library, 10 West 35th Street, Chicago, Illinois 60616 (1)

OTHER (continued)

- Johns Hopkins University, Applied Physics Laboratory, ATTN: Document Library, 8621 Georgia Avenue, Silver Spring, Maryland 20910 (1)
- Dr. R. F. Kallman, Department of Radiology, Stanford University, Palo Alto, California 94305 (1)
- Dr. L. S. Kelly, Donner Laboratory, University of California at Berkeley, Berkeley, California 94720 (1)
- Los Alamos Scientific Laboratory, ATTN: Report Librarian, P. O. Box 1663, Los Alamos, New Mexico 87544 (1)
- Director, Nuclear Science Center, Louisiana State University, Baton Rouge, Louisiana 70803 (2)
- Lovelace Foundation for Medical Education & Research, Document Library, 5200 Gibson Boulevard, S. E., Albuquerque, New Mexico 87108 (1)
- Dr. Ross A. McFarland, Guggenheim Professor of Aerospace Health & Safety, Harvard School of Public Health, 665 Huntington Avenue, Boston, Massachusetts 02115 (1)
- Dr. J. I. Marcum, Rand Corporation, 1700 Main Street, Santa Monica, California 90401 (1)
- Massachusetts Institute of Technology, M.I.T. Libraries, Technical Reports, Room 14 E-210, Cambridge, Massachusetts 02139 (1)
- Dr. Charles W. Mays, Physics Group Leader, Radiobiology Division, University of Utah, Salt Lake City, Utah 84112 (1)
- Dr. B. D. Newsom, Colony Oaks, Apt. 32, 18100 Nassau Bay Drive, Nassau Bay, Texas 77058 (1)
- Ohio State University, Nuclear Reactor Laboratory, 1298 Kinnear Road, Columbus, Ohio 43212 (1)
- Dr. Harvey M. Patt, Laboratory of Radiobiology, University of California, San Francisco Medical Center, San Francisco, California 94122 (1)
- Purdue University, Nuclear Engineering Library, Lafayette, Indiana 47907 (1)
- Dr. S. M. Reichard, Director, Division of Radiobiology, Medical College of Georgia, Augusta, Georgia 30902 (1)
- University of Rochester, Atomic Energy Project Library, P. O. Box 287, Station 3, Rochester, New York 14620 (1)
- Dr. H. H. Rossi, 630 West 168th Street, New York, New York 10032 (1)
- Dr. Eugene L. Saenger, Director, Radioisotope Laboratory, Cincinnati General Hospital, Cincinnati, Ohio 45229 (1)
- Sandia Corporation Library, P. O. Box 5800, Albuquerque, New Mexico 87115 (1)
- Scientific Committee on the Effects of Atomic Radiation, ATTN: Library, United Nations Room 3267, United Nations Plaza, New York, New York 10017 (1)
- Scope Publications, Franklin Station, P. O. Box 7407, Washington, D. C. 20004 (1)
- University of Southern California, Nuclear Physics Laboratory, University Park, Los Angeles, California 90007 (1)
- Dr. Arthur R. Tamplin, Biophysicist, Information Integration Group, University of California, Lawrence Radiation Laboratory, L-612, Livermore, California 94550 (1)
- Radiation Biology Laboratory, Texas Engineering Experiment Station, Texas A. & M. University, College Station, Texas 77840 (2)
- Texas Nuclear Corporation, ATTN: Director of Research, Box 9267 Allandale Station, Austin, Texas 78756 (1)
- Western Reserve University, Department of Radiology, Division of Radiation Biology, Cleveland, Ohio 44106 (1)
- Mr. Lionel Zamore, 601 Brightwater Court, Brooklyn, New York 11235 (1)

FOREIGN

- International Atomic Energy Agency, Kaerntnerring 11, Vienna I. 1010, Austria (1)
- European Atomic Energy Community, C.E.E.A., Library, 51 rue Belliard, Brussels 4, Belgium (1)
- Dr. L. G. Lajtha, Paterson Laboratories, Christie Hospital and Holt Radium Institute, Manchester, England (1)
- Dr. L. F. Lamerton, Biophysics Department, Institute of Cancer Research, Surrey Branch, Belmont, Sutton, Surrey, England (1)
- National Lending Library for Science and Technology, Boston Spa, Yorkshire, England (1)
- Directorate of Medical and Health Services, FAF (Federal Armed Forces), Bonn, Ermekeilstr. 27, West Germany (1)
- Abteilung fur Strahlenbiologie im Institut fur Biophysik der Universitat Bonn, 53 Bonn-Venusberg, Annaberger Weg 15, Federal Republic of Germany (2)
- Prof. Dr. H. Langendorff, Direktor des Radiologischen Instituts der Universitat, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Dr. Helmut Mitschrich, Akademie des Sanitaets-und Gesundheits, Weseus BW, Spezialstab ATV, 8 Muenchen Schwere-Reiterstr. 4, Germany (2)
- Prof. Dr. F. Wachsmann, Gesellschaft fur Strahlenforschung m.b.H., 8042 Neuherberg bei Muenchen, Institut fur Strahlenschutz, Ingolstadter Landstrasse 1, Muenchen, Germany (1)
- Dr. M. Feldman, Section of Cell Biology, The Weizmann Institute of Science, Rehovoth, Israel (1)
- Dr. G. W. Barendsen, Radiobiological Institute TNO, Rijswijk, Netherlands (1)
- Dr. L. M. van Putten, Radiobiological Institute TNO, 151 Lance Kleiweg, Rijswijk 2 H., Netherlands (1)
- Puerto Rico Nuclear Center, ATTN: Reading Room, College Station, Mayaguez, Puerto Rico 00708 (2)
- Dr. H. Cottier, Pathological Institut der Universitat, Bern, Switzerland (1)

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Armed Forces Radiobiology Research Institute Defense Atomic Support Agency Bethesda, Maryland 20014		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP N/A	
3. REPORT TITLE EFFECT OF IONIZING RADIATIONS ON DISTRIBUTION OF PLASMA PROTEIN-BOUND NEUTRAL HEXOSES IN MICE AND DOGS			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) Evans, A. S.			
6. REPORT DATE December 1969	7a. TOTAL NO. OF PAGES 20	7b. NO. OF REFS 10	
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) AFRRI SR69-24		
b. PROJECT NO.			
c. MC 3 90102	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
d.			
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Defense Atomic Support Agency Washington, D. C. 20305	
13. ABSTRACT Comparison of changes in concentration of protein-bound carbohydrates (PBC) as neutral hexoses in the plasma of C3H mice and beagles as a function of time after exposure to mixed gamma-neutron radiations revealed that the two species react similarly. The animals which died exhibited a marked increase in plasma PBC concentration while the survivors deviated only slightly from their preirradiation values. Analytical acrylamide gel electrophoresis demonstrated that distinctive, progressive postirradiation changes occurred in the plasma of the animals which succumbed. The most striking changes were in the components tentatively identified as transferrin, haptoglobins, and α_2 -macroglobulin. Preparative fractionation of plasma of both mice and dogs by combined molecular sieving and ion exchange chromatography revealed that, in the four fractions uniformly obtained, absolute changes in PBC regularly occurred only in the first and third. The protein composition of these fractions and their significance as a prognostic tool in radiation injury are discussed. The suggestion that the preirradiation PBC concentration may constitute a crude index to radiosensitivity was confirmed under conditions specifically designed to test the hypothesis.			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.